



Active substances set

Search phrase: ramucirumab

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

Malignant stomach cancer

	Ramucirumab in combination with paclitaxel is indicated for		
	the treatment of adult patients with advanced gastric		
	cancer or gastro-oesophageal junction adenocarcinoma		
	with disease progression after prior platinum and		
	fluoropyrimidine chemotherapy. Ramucirumab	0	REIMBURSEMENT
Ramucirumab	monotherapy is indicated for the treatment of adult		WITH RESTRICTIONS
	patients with advanced gastric cancer or gastro-		ESMO
	oesophageal junction adenocarcinoma with disease		
	progression after prior platinum or fluoropyrimidine		
	chemotherapy, for whom treatment in combination with		
	paclitaxel is not appropriate.		

Colon and rectum cancer

Ramucirumab, in combination with FOLFIRI (irinotecan, folinic acid, and 5-fluorouracil), is indicated for the treatment of adult patients with metastatic colorectal cancer (mCRC) with disease progression on or after prior therapy with bevacizumab, oxaliplatin and a fluoropyrimidine.

NO REIMBURSEMENT ESMO

Liver cancer

Ramucirumab

Ramucirumab monotherapy is indicated for the treatment of adult patients with advanced or unresectable hepatocellular carcinoma who have a serum alpha fetoprotein (AFP) of \geq 400 ng/ml and who have been previously treated with sorafenib.



NO REIMBURSEMENT

ESMO

Tracheal, bronchus, and lung cancer

Ramucirumab in combination with erlotinib is indicated for the first-line treatment of adult patients with metastatic non-small cell lung cancer with activating epidermal growth factor receptor (EGFR) mutations. Ramucirumab in combination with docetaxel is indicated for the treatment of adult patients with locallyadvanced or metastatic non-small cell lung cancer with disease progression after platinumbased chemotherapy.

Ramucirumab

Ramucirumab

adult patients with loc cell lung cancer with o based chemotherapy.