



Active substances set

Search phrase: trifluridine / tipiracil hydrochloride

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

Malignant stomach cancer

		Trifluridine / Tipiracil Hydrochloride is indicated as		
	Trifluridine /	monotherapy for the treatment of adult patients with		REIMBURSEMENT
	tipiracil	metastatic gastric cancer including adenocarcinoma of the		WITH RESTRICTIONS
		gastroesophageal junction, who have been previously		
	hydrochloride	treated with at least two prior systemic treatment regimens	\checkmark	ESMO
		for advanced disease.		

Colon and rectum cancer

	Trifluridine / Tipiracil Hydrochloride is indicated in	
	combination with bevacizumab for the treatment of adult	
	patients with metastatic colorectal cancer (CRC) who have	
	received two prior anticancer treatment regimens including	
Trifluridine /	fluoropyrimidine-, oxaliplatin- and irinotecan-based	
	chemotherapies, anti-VEGF agents, and/or anti-EGFR	FULL
tipiracil	agents. Trifluridine / Tipiracil Hydrochloride is indicated as	REIMBURSEMENT
hydrochloride	monotherapy for the treatment of adult patients with	ESMO
	metastatic colorectal cancer who have been previously	
	treated with, or are not considered candidates for, available	
	therapies including fluoropyrimidine-, oxaliplatin- and	
	irinotecan-based chemotherapies, anti-VEGF agents, and	
	anti-EGFR agents.	