

## Active substances set

**Search phrase:** blinatumomab

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

### Acute lymphoid leukemia

Blinatumomab is indicated as monotherapy for the treatment of adults with CD19 positive relapsed or refractory B-precursor acute lymphoblastic leukaemia (ALL). Patients with Philadelphia chromosome positive B-precursor ALL should have failed treatment with at least 2 tyrosine kinase inhibitors (TKIs) and have no alternative treatment options. Blinatumomab is indicated as monotherapy for the treatment of adults with Philadelphia chromosome negative CD19 positive B-precursor ALL in first or second complete remission with minimal residual disease (MRD) greater than or equal to 0.1%.

Blinatumomab is indicated as monotherapy for the treatment of paediatric patients aged 1 year or older with Philadelphia chromosome negative CD19 positive B-precursor ALL which is refractory or in relapse after receiving at least two prior therapies or in relapse after receiving prior allogeneic haematopoietic stem cell transplantation. Blinatumomab is indicated as monotherapy for the treatment of paediatric patients aged 1 year or older with high-risk first relapsed Philadelphia chromosome negative CD19 positive B-precursor ALL as part of the consolidation therapy. Blinatumomab is indicated as monotherapy as part of consolidation therapy for the treatment of adult patients with newly diagnosed Philadelphia chromosome negative CD19 positive B-cell precursor ALL.

#### Blinatumomab

 **REIMBURSEMENT  
WITH RESTRICTIONS**

 **ESMO**