# Active substances set

**Search phrase:** Malignant breast cancer

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

## Malignant breast cancer

| | | |
|---|---|---|
| **Talazoparib** | Talazoparib is indicated as monotherapy for the treatment of adult patients with germline BRCA1/2-mutations, who have HER2-negative locally advanced or metastatic breast cancer. Patients should have been previously treated with an anthracycline and/or a taxane in the (neo)adjuvant, locally advanced or metastatic setting unless patients were not suitable for these treatments. Patients with hormone receptor (HR)-positive breast cancer should have been treated with a prior endocrine-based therapy, or be considered unsuitable for endocrine-based therapy. | NO REIMBURSEMENT<br>ESMO |
| **Capivasertib** | Capivasertib is indicated in combination with fulvestrant for the treatment of adult patients with oestrogen receptor (ER)-positive, HER2-negative locally advanced or metastatic breast cancer with one or more PIK3CA/AKT1/PTEN-alterations following recurrence or progression on or after an endocrine-based regimen. In pre- or perimenopausal women, Capivasertib plus fulvestrant should be combined with a luteinising hormone releasing hormone (LHRH) agonist. For men, administration of LHRH agonist according to current clinical practice standards should be considered. | NO REIMBURSEMENT<br>ESMO |
| **Datopotamab deruxtecan** | Breast cancer Datopotamab deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-negative breast cancer who have received endocrine therapy and at least one line of chemotherapy in the advanced setting. | NO REIMBURSEMENT<br>ESMO |

| | | |
|---|---|---|
| **Imlunestrant** | Imlunestrant is indicated as monotherapy for the treatment of adult patients with oestrogen receptor (ER)-positive, HER2-negative, locally advanced or metastatic breast cancer with an activating ESR1-mutation, who have disease progression following prior treatment with an endocrine based regimen. | NO REIMBURSEMENT<br>ESMO |
| **Eribulin** | Eribulin is indicated for the treatment of adult patients with locally advanced or metastatic breast cancer who have progressed after at least one chemotherapeutic regimen for advanced disease. Prior therapy should have included an anthracycline and a taxane in either the adjuvant or metastatic setting unless patients were not suitable for these treatments. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Everolimus** | Everolimus is indicated for the treatment of hormone receptor-positive, HER2/neu negative advanced breast cancer, in combination with exemestane, in postmenopausal women without symptomatic visceral disease after recurrence or progression following a non-steroidal aromatase inhibitor. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Pertuzumab** | Early breast cancer Pertuzumab is indicated for use in combination with trastuzumab and chemotherapy in: - the neoadjuvant treatment of adult patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer at high risk of recurrence; - the adjuvant treatment of adult patients with HER2-positive early breast cancer at high risk of recurrence. Metastatic breast cancer Pertuzumab is indicated for use in combination with trastuzumab and docetaxel in adult patients with HER2-positive metastatic or locally recurrent unresectable breast cancer, who have not received previous anti-HER2 therapy or chemotherapy for their metastatic disease. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Palbociclib** | Palbociclib is indicated for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer: - in combination with an aromatase inhibitor; - in combination with fulvestrant in women who have received prior endocrine therapy. In pre- or perimenopausal women, the endocrine therapy should be combined with a luteinizing hormone-releasing hormone (LHRH) agonist. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Atezolizumab** | Atezolizumab in combination with nab-paclitaxel is indicated for the treatment of adult patients with unresectable locally advanced or metastatic triple-negative breast cancer (TNBC) whose tumours have PD-L1 expression ≥ 1% and who have not received prior chemotherapy for metastatic disease. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |

| | | |
|---|---|---|
| **Neratinib** | Neratinib is indicated for the extended adjuvant treatment of adult patients with early-stage hormone receptor positive HER2-overexpressed/amplified breast cancer and who completed adjuvant trastuzumab-based therapy less than one year ago. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Olaparib** | Olaparib is indicated as: - monotherapy or in combination with endocrine therapy for the adjuvant treatment of adult patients with germline BRCA1/2-mutations who have HER2-negative, high risk early breast cancer previously treated with neoadjuvant or adjuvant chemotherapy. - monotherapy for the treatment of adult patients with germline BRCA1/2-mutations, who have HER2 negative locally advanced or metastatic breast cancer. Patients should have previously been treated with an anthracycline and a taxane in the (neo)adjuvant or metastatic setting unless patients were not suitable for these treatments. Patients with hormone receptor (HR)-positive breast cancer should also have progressed on or after prior endocrine therapy, or be considered unsuitable for endocrine therapy. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Alpelisib** | Alpelisib is indicated in combination with fulvestrant for the treatment of postmenopausal women, and men, with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with a PIK3CA mutation after disease progression following endocrine therapy as monotherapy. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Pertuzumab / trastuzumab** | Early breast cancer (EBC) Pertuzumab / trastuzumab is indicated for use in combination with chemotherapy in: - the neoadjuvant treatment of adult patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer at high risk of recurrence, - the adjuvant treatment of adult patients with HER2-positive early breast cancer at high risk of recurrence. Metastatic breast cancer (MBC) Pertuzumab / trastuzumab is indicated for use in combination with docetaxel in adult patients with HER2-positive metastatic or locally recurrent unresectable breast cancer, who have not received previous anti-HER2 therapy or chemotherapy for their metastatic disease. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |

| | | |
|---|---|---|
| **Trastuzumab deruxtecan** | Trastuzumab Deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received one or more prior anti-HER2-based regimens. Trastuzumab Deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-low breast cancer who have received prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Tucatinib** | Tucatinib is indicated in combination with trastuzumab and capecitabine for the treatment of adult patients with HER2-positive locally advanced or metastatic breast cancer who have received at least 2 prior anti-HER2 treatment regimens. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Pembrolizumab** | Pembrolizumab, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early-stage triple-negative breast cancer (TNBC) at high risk of recurrence. Pembrolizumab, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple-negative breast cancer in adults whose tumours express PD-L1 with a CPS ≥ 10 and who have not received prior chemotherapy for metastatic disease. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Sacituzumab govitecan** | Sacituzumab Govitecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic triple-negative breast cancer (mTNBC) who have received two or more prior systemic therapies, including at least one of them for advanced disease. Sacituzumab Govitecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-negative breast cancer who have received endocrine-based therapy, and at least two additional systemic therapies in the advanced setting. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |
| **Elacestrant** | Elacestrant monotherapy is indicated for the treatment of postmenopausal women, and men, with estrogen receptor (ER)-positive, HER2-negative, locally advanced or metastatic breast cancer with an activating ESR1 mutation who have disease progression following at least one line of endocrine therapy including a CDK 4/6 inhibitor. | REIMBURSEMENT WITH RESTRICTIONS<br>ESMO |

| | | |
|---|---|---|
| **Trastuzumab emtansine** | Trastuzumab Emtansine, as a single agent, is indicated for the adjuvant treatment of adult patients with HER2-positive early breast cancer who have residual invasive disease, in the breast and/or lymph nodes, after neoadjuvant taxane-based and HER2-targeted therapy. Trastuzumab Emtansine, as a single agent, is indicated for the treatment of adult patients with HER2-positive, unresectable locally advanced or metastatic breast cancer who previously received trastuzumab and a taxane, separately or in combination. Patients should have either: - Received prior therapy for locally advanced or metastatic disease, or - Developed disease recurrence during or within six months of completing adjuvant therapy. | FULL REIMBURSEMENT<br>ESMO |
| **Ribociclib** | Early breast cancer Ribociclib in combination with an aromatase inhibitor is indicated for the adjuvant treatment of patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative early breast cancer at high risk of recurrence (see section 5.1 for selection criteria). In pre- or perimenopausal women, or in men, the aromatase inhibitor should be combined with a luteinising hormone-releasing hormone (LHRH) agonist. Advanced or metastatic breast cancer Ribociclib is indicated for the treatment of women with HR-positive, HER2-negative locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulvestrant as initial endocrine-based therapy, or in women who have received prior endocrine therapy. In pre- or perimenopausal women, the endocrine therapy should be combined with a LHRH agonist. | FULL REIMBURSEMENT<br>ESMO |
| **Abemaciclib** | Early breast cancer Abemaciclib in combination with endocrine therapy is indicated for the adjuvant treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, node-positive early breast cancer at high risk of recurrence. In pre- or perimenopausal women, aromatase inhibitor endocrine therapy should be combined with a luteinising hormone-releasing hormone (LHRH) agonist. Advanced or metastatic breast cancer Abemaciclib is indicated for the treatment of women with hormone receptor (HR) positive, human epidermal growth factor receptor 2 (HER2) negative locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulvestrant as initial endocrine-based therapy, or in women who have received prior endocrine therapy. In pre- or perimenopausal women, the endocrine therapy should be combined with a LHRH agonist. | FULL REIMBURSEMENT<br>ESMO |

| | | |
|---|---|---|
| **Inavolisib** | Inavolisib, in combination with palbociclib and fulvestrant, is indicated for the treatment of adult patients with PIK3CA-mutated, oestrogen receptor (ER)-positive, HER2-negative, locally advanced or metastatic breast cancer, following recurrence on or within 12 months of completing adjuvant endocrine treatment. Patients previously treated with a CDK 4/6 inhibitor in the (neo)adjuvant setting should have had an interval of at least 12 months between termination of CDK 4/6 inhibitor treatment and the detection of recurrence. In pre/perimenopausal women and in men, endocrine therapy should be combined with a luteinising hormone-releasing hormone (LHRH) agonist. | FULL REIMBURSEMENT<br>ESMO |