



Active substances set

Search phrase: Malignant breast cancer

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

Malignant breast cancer

Talazoparib is indicated as monotherapy for the treatment of adult patients with germline BRCA1/2-mutations, who have HER2-negative locally advanced or metastatic breast cancer. Patients should have been previously treated with an anthracycline and/or a taxane in the (neo)adjuvant, locally advanced or metastatic setting unless patients were not suitable for these treatments. Patients with hormone receptor (HR)-positive breast cancer should have been treated with a prior endocrine-based therapy, or be considered unsuitable for endocrine-based therapy.





Elacestrant

Talazoparib

Elacestrant monotherapy is indicated for the treatment of postmenopausal women, and men, with estrogen receptor (ER)-positive, HER2-negative, locally advanced or metastatic breast cancer with an activating ESR1 mutation who have disease progression following at least one line of endocrine therapy including a CDK 4/6 inhibitor.

Capivasertib is indicated in combination with fulvestrant for





Capivasertib

the treatment of adult patients with oestrogen receptor (ER)-positive, HER2-negative locally advanced or metastatic breast cancer with one or more PIK3CA/AKT1/PTENalterations following recurrence or progression on or after an endocrine-based regimen. In pre- or perimenopausal women, Capivasertib plus fulvestrant should be combined with a luteinising hormone releasing hormone (LHRH) agonist. For men, administration of LHRH agonist according to current clinical practice standards should be considered.





Datopotamab deruxtecan

Breast cancer Datopotamab deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-negative breast cancer who have received endocrine therapy and at least one line of chemotherapy in the advanced setting.

Inavolisib, in combination with palbociclib and fulvestrant.





Inavolisib

is indicated for the treatment of adult patients with PIK3CA-mutated, oestrogen receptor (ER)-positive, HER2-negative, locally advanced or metastatic breast cancer, following recurrence on or within 12 months of completing adjuvant endocrine treatment. Patients previously treated with a CDK 4/6 inhibitor in the (neo)adjuvant setting should have had an interval of at least 12 months between termination of CDK 4/6 inhibitor treatment and the detection of recurrence. In pre/perimenopausal women and in men, endocrine therapy should be combined with a luteinising hormone-releasing hormone (LHRH) agonist.





Eribulin

Eribulin is indicated for the treatment of adult patients with locally advanced or metastatic breast cancer who have progressed after at least one chemotherapeutic regimen for advanced disease. Prior therapy should have included an anthracycline and a taxane in either the adjuvant or metastatic setting unless patients were not suitable for these treatments.

- REIMBURSEMENT
 WITH RESTRICTIONS
- S ESMO

Everolimus

Everolimus is indicated for the treatment of hormone receptor-positive, HER2/neu negative advanced breast cancer, in combination with exemestane, in postmenopausal women without symptomatic visceral disease after recurrence or progression following a non-steroidal aromatase inhibitor.

- REIMBURSEMENT WITH RESTRICTIONS
- **ESMO**

Pertuzumab

Early breast cancer Pertuzumab is indicated for use in combination with trastuzumab and chemotherapy in: - the neoadjuvant treatment of adult patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer at high risk of recurrence; - the adjuvant treatment of adult patients with HER2-positive early breast cancer at high risk of recurrence. Metastatic breast cancer Pertuzumab is indicated for use in combination with trastuzumab and docetaxel in adult patients with HER2-positive metastatic or locally recurrent unresectable breast cancer, who have not received previous anti-HER2 therapy or chemotherapy for their metastatic disease.

- REIMBURSEMENT WITH RESTRICTIONS
- S ESMO

Palbociclib

Palbociclib is indicated for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer: - in combination with an aromatase inhibitor; - in combination with fulvestrant in women who have received prior endocrine therapy. In pre- or perimenopausal women, the endocrine therapy should be combined with a luteinizing hormone-releasing hormone (LHRH) agonist.

- REIMBURSEMENT
 WITH RESTRICTIONS
 - **ESMO**

Neratinib

Neratinib is indicated for the extended adjuvant treatment of adult patients with early-stage hormone receptor positive HER2-overexpressed/amplified breast cancer and who completed adjuvant trastuzumab-based therapy less than one year ago.

- REIMBURSEMENT
 WITH RESTRICTIONS
- **ESMO**

Olaparib

Olaparib is indicated as: - monotherapy or in combination with endocrine therapy for the adjuvant treatment of adult patients with germline BRCA1/2-mutations who have HER2-negative, high risk early breast cancer previously treated with neoadjuvant or adjuvant chemotherapy. - monotherapy for the treatment of adult patients with germline BRCA1/2-mutations, who have HER2 negative locally advanced or metastatic breast cancer. Patients should have previously been treated with an anthracycline and a taxane in the (neo)adjuvant or metastatic setting unless patients were not suitable for these treatments. Patients with hormone receptor (HR)-positive breast cancer should also have progressed on or after prior endocrine therapy, or be considered unsuitable for endocrine therapy.

- REIMBURSEMENT
 WITH RESTRICTIONS
 - S ESMO

Alpelisib

Alpelisib is indicated in combination with fulvestrant for the treatment of postmenopausal women, and men, with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with a PIK3CA mutation after disease progression following endocrine therapy as monotherapy.

- REIMBURSEMENT
 WITH RESTRICTIONS
- **ESMO**

Trastuzumab deruxtecan

Trastuzumab Deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received one or more prior anti-HER2-based regimens. Trastuzumab Deruxtecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-low breast cancer who have received prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy.

- REIMBURSEMENT
 WITH RESTRICTIONS
- **ESMO**

Tucatinib

Tucatinib is indicated in combination with trastuzumab and capecitabine for the treatment of adult patients with HER2-positive locally advanced or metastatic breast cancer who have received at least 2 prior anti-HER2 treatment regimens.





Sacituzumab govitecan

Sacituzumab Govitecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic triple-negative breast cancer (mTNBC) who have received two or more prior systemic therapies, including at least one of them for advanced disease. Sacituzumab Govitecan as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-negative breast cancer who have received endocrine-based therapy, and at least two additional systemic therapies in the advanced setting.

Trastuzumab Emtansine, as a single agent, is indicated for

- REIMBURSEMENT
 WITH RESTRICTIONS
- S ESMO

Trastuzumab emtansine

the adjuvant treatment of adult patients with HER2-positive early breast cancer who have residual invasive disease, in the breast and/or lymph nodes, after neoadjuvant taxane-based and HER2-targeted therapy. Trastuzumab Emtansine, as a single agent, is indicated for the treatment of adult patients with HER2-positive, unresectable locally advanced or metastatic breast cancer who previously received trastuzumab and a taxane, separately or in combination. Patients should have either: - Received prior therapy for locally advanced or metastatic disease, or - Developed disease recurrence during or within six months of completing adjuvant therapy.



ESMO

Ribociclib

Early breast cancer Ribociclib in combination with an aromatase inhibitor is indicated for the adjuvant treatment of patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative early breast cancer at high risk of recurrence (see section 5.1 for selection criteria). In pre- or perimenopausal women, or in men, the aromatase inhibitor should be combined with a luteinising hormone-releasing hormone (LHRH) agonist. Advanced or metastatic breast cancer Ribociclib is indicated for the treatment of women with HR-positive, HER2-negative locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulvestrant as initial endocrine-based therapy, or in women who have received prior endocrine therapy. In pre- or perimenopausal women, the endocrine therapy should be combined with a LHRH agonist.



ESMO

Atezolizumab

Abemaciclib

Atezolizumab in combination with nab-paclitaxel is indicated for the treatment of adult patients with unresectable locally advanced or metastatic triple-negative breast cancer (TNBC) whose tumours have PD-L1 expression ≥ 1% and who have not received prior chemotherapy for metastatic disease.





Early breast cancer Abemaciclib in combination with endocrine therapy is indicated for the adjuvant treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, nodepositive early breast cancer at high risk of recurrence. In pre- or perimenopausal women, aromatase inhibitor endocrine therapy should be combined with a luteinising hormone-releasing hormone (LHRH) agonist. Advanced or metastatic breast cancer Abemaciclib is indicated for the

treatment of women with hormone receptor (HR) positive, human epidermal growth factor receptor 2 (HER2) negative locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulvestrant as initial endocrine-based therapy, or in women who have received prior endocrine therapy. In pre- or perimenopausal





Pembrolizumab, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early-stage triple-negative breast cancer (TNBC) at high Pembrolizumab risk of recurrence. Pembrolizumab, in combination with

women, the endocrine therapy should be combined with a

LHRH agonist.

chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple-negative breast cancer in adults whose tumours express PD-L1 with a CPS \geq 10 and who have not received prior chemotherapy for metastatic disease.



