

## Active substances set

**Search phrase:** Malignant stomach cancer

Below you will find a list of active substances registered by the European Medical Agency (EMA) in the last 15 years, recommended by the European Society of Clinical Oncology (ESMO) and their reimbursement status in the country.

### Malignant stomach cancer

#### **Trifluridine / tipiracil hydrochloride**

Trifluridine / Tipiracil Hydrochloride is indicated as monotherapy for the treatment of adult patients with metastatic gastric cancer including adenocarcinoma of the gastroesophageal junction, who have been previously treated with at least two prior systemic treatment regimens for advanced disease.

 **NO REIMBURSEMENT**  
 **ESMO**

#### **Trastuzumab deruxtecan**

Trastuzumab Deruxtecan as monotherapy is indicated for the treatment of adult patients with advanced HER2-positive gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen.

 **NO REIMBURSEMENT**  
 **ESMO**

#### **Zolbetuximab**

Zolbetuximab, in combination with fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the first-line treatment of adult patients with locally advanced unresectable or metastatic HER2-negative gastric or gastroesophageal junction (GEJ) adenocarcinoma whose tumours are Claudin (CLDN) 18.2 positive.

 **NO REIMBURSEMENT**  
 **ESMO**

## Pembrolizumab

Pembrolizumab as monotherapy is indicated for the treatment of the following microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) tumours in adults with unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy. Pembrolizumab, in combination with trastuzumab, fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction (GEJ) adenocarcinoma in adults whose tumours express PD-L1 with a CPS  $\geq 1$ . Pembrolizumab, in combination with fluoropyrimidine and platinum -containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2 - negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS  $\geq 1$ .

 **REIMBURSEMENT WITH RESTRICTIONS**

 **ESMO**

## Ramucirumab

Ramucirumab in combination with paclitaxel is indicated for the treatment of adult patients with advanced gastric cancer or gastro-oesophageal junction adenocarcinoma with disease progression after prior platinum and fluoropyrimidine chemotherapy. Ramucirumab monotherapy is indicated for the treatment of adult patients with advanced gastric cancer or gastro-oesophageal junction adenocarcinoma with disease progression after prior platinum or fluoropyrimidine chemotherapy, for whom treatment in combination with paclitaxel is not appropriate.

 **FULL REIMBURSEMENT**

 **ESMO**

## Nivolumab

Nivolumab in combination with fluoropyrimidine- and platinum-based combination chemotherapy is indicated for the first-line treatment of adult patients with HER2-negative advanced or metastatic gastric, gastro-oesophageal junction (GEJ) or oesophageal adenocarcinoma whose tumours express PD-L1 with a combined positive score (CPS)  $\geq 5$ .

 **FULL REIMBURSEMENT**

 **ESMO**